

Understanding the development of psychopathy: progress and challenges

E. Viding* and E. J. McCrory

Division of Psychology and Language Sciences, University College London, Gower St, London, WC1E 6BT, UK

Psychopathy is an adult condition that incurs substantial societal and individual costs. Here we review neurocognitive and genetically informative studies that shed light on how and why this condition emerges. Children cannot present with psychopathy. However, the presence of callous–unemotional (CU) traits can distinguish a group of children who are at elevated risk of psychopathy in adulthood. These children display diminished empathy and guilt and show attenuated brain activation to distress cues in others. Genetically informative studies indicate that individual differences in CU traits show moderate-to-strong heritability, but that protective environmental factors can counter heritable risk. On the basis of the extant research findings, we speculate on what might represent the priorities for research over the next decade. We also consider the clinical implications of these research findings. In particular, we consider the importance of delineating what precisely works for children with CU traits (and their parents) and the ways in which intervention and prevention programs may be optimized to improve engagement as well as clinical outcomes.

Received 10 April 2017; Revised 25 August 2017; Accepted 29 August 2017; First published online 16 October 2017

Key words: Antisocial, brain, callous-unemotional, development, empathy, genetic, psychopathy.

Background

Psychopathy is a personality disorder characterized by lack of empathy and guilt, shallow affect, manipulation of other people, and severe, premeditated, and violent antisocial behavior (Hare & Neumann, 2008). Although individuals with criminal psychopathy share many features, including impulsivity, with other antisocial offenders, what really sets them apart is a relative absence of guilt for their actions, little concern for the suffering of their fellow human beings, shallow affiliative bonds, and a tendency to not show loyalty unless it is in their own interest (Hare & Neumann, 2008; Kiehl, 2015). Individuals with psychopathy incur substantial societal costs, both as a direct financial consequence of their offending behavior and lack of normal participation in working life, but also in terms of the emotional and psychological costs to their victims. Given the serious nature of the disorder, researchers and clinicians are motivated to understand how psychopathy develops and how it might be prevented.

Measuring psychopathy in adults

Diagnostic manuals such as the *Diagnostic and Statistical Manual of Mental Disorders Fifth Edition* (DSM V) recognize

antisocial personality disorder (ASPD), which is characterized by aggression, impulsivity, and violation of the rights of others. However, there is no specific diagnostic category of psychopathy. Only a minority of individuals diagnosed with ASPD meet criteria for psychopathy as diagnosed by the Psychopathy Checklist – Revised (Hare, 2003), a standard instrument used in criminal justice settings that measures affective, interpersonal, lifestyle, and antisocial features associated with psychopathic personality. Other instruments such as the Triarchic Psychopathy Measure (Patrick, 2010), the Self-Report Psychopathy scale (Paulhus *et al.* 2016), and the Psychopathic Personality Inventory (Lilienfeld & Andrews, 1996) are also used to reliably measure psychopathic traits in a variety of settings. While there are differences between these measures, all capture aspects of callous presentation, lack of empathy, impulsivity, and disinhibition. Some, such as the Triarchic Psychopathy Measure and the Psychopathic Personality Inventory, also measure additional domains such as boldness/fearlessness.

Conceptualizing psychopathy within a developmental context

Can a child present with psychopathy? This remains a complex and controversial question. In our view, it is inappropriate (and inaccurate) to label any child as a psychopath. The label of psychopathy is frequently (but not always) used in the context of intractable criminality and violent behavior. Although the data are clear that

* Address for correspondence: E. Viding, Division of Psychology and Language Sciences, University College London, Gower St, London, WC1E 6BT, UK.
(Email: e.viding@ucl.ac.uk)

comparable personality features and associated neurocognitive indicators can be identified in both adults with psychopathy and children with psychopathic features (Blair *et al.* 2014; Viding & McCrory, 2015), we also know that these features and associated behaviors are not fixed during childhood (e.g. Fontaine *et al.* 2010; Waller *et al.* 2013). In this light, the negative connotations of a psychopathy label seem overly harsh in reference to children, especially as we know that developmental outcomes are not certain. This is the reason that we prefer to talk about developmental risk for psychopathy or psychopathic traits/features when we discuss findings relating to children. Our position is also in line with the fact that personality disorder diagnoses more broadly are reserved for adult populations.

However, while psychopathic features, in common with other personality features, may not be fixed across childhood, an established body of research now demonstrates that they are predictive of increased risk for persistent antisocial behavior and adult psychopathic presentation (Lynam *et al.* 2007; Viding & McCrory, 2015). Different measures, such as the Antisocial Process Screening Device (Frick & Hare, 2001), Youth Psychopathic Traits Inventory (Andershed *et al.* 2002), Inventory of Callous Unemotional Traits (Essau *et al.* 2006), and Child Problematic Traits Inventory (Colins *et al.* 2014) reliably capture psychopathic personality traits including callous, unempathetic, interpersonal, and impulsive features. In combination, these traits substantially increase the risk of persistent antisocial behavior in children and youth (Frick *et al.* 2014; Viding & McCrory, 2015; Frogner *et al.* 2016). The diagnostic schemes for conduct disorder capture many of the impulsive and interpersonal aspects associated with psychopathic presentation (some of which are shared with other individuals with antisocial behavior). The recent addition to the American Psychiatric Association DSM 5 (American Psychiatric Association, 2013) of a 'limited prosocial emotions' specifier [or callous-unemotional (CU) traits as they have been termed in research contexts] now also provides some coverage of the affective/unempathetic features of psychopathy. In line with the DSM 5 approach, many researchers, ourselves included, have used a measure of CU traits alongside a measure disruptive behaviors (capturing impulsive and interpersonal characteristics) to differentiate a group of children with disruptive behaviors who are at significantly elevated risk for persistent antisocial behavior/psychopathy, but who differ from other children with disruptive behavior in their affective presentation.

The current review

In this review, we will provide a short overview of *neurocognitive* studies that have focused on children with

disruptive behaviors and CU traits. Findings from these studies can help us understand the degree to which charting disruptive behaviors together with CU traits identifies a group of children who are comparable to adults who present with psychopathy. We will also discuss *genetically informative studies* that have focused on psychopathic personality features (often CU traits) or on disruptive behaviors in the presence *v.* absence of such features. We have chosen to focus this review on the neurocognitive and genetically informative findings for two reasons. First, we think that findings from these two neurobiological areas can inform each other and a possible 'roadmap' for future research efforts that can bring us closer to understanding how psychopathy develops. Second, the extant research findings have a number of clear clinical implications, which we will also discuss.

Findings from neurocognitive studies

Behaviorally, children with disruptive behaviors and CU traits show a marked lack of empathy or guilt. They often engage in proactive, instrumental aggression, seem impervious to sanctions, and do not appear to exhibit the affiliative needs and goals that characterize typical children (Frick *et al.* 2014; Viding & McCrory, 2015). Given this profile, many of the experimental studies on children with disruptive behaviors and CU traits have focused on how they process emotions, whether they empathize with others, how they attend to caregivers, and whether they change their behavior following punishment (see Kimonis *et al.* 2008; Marsh & Blair, 2008; Jones *et al.* 2010; Dadds *et al.* 2011, 2012; Sylvers *et al.* 2011; de Wied *et al.* 2012; Schwenck *et al.* 2012; Blair *et al.* 2014; Hodsoll *et al.* 2014; Bedford *et al.* 2015; Martin-Key *et al.* 2016; White *et al.* 2016a, b). The majority of these studies have documented that, compared with typically developing children, those with disruptive behaviors and CU traits are less likely to attend to, react to, and recognize affective stimuli, including distress cues such as fearful and sad expressions of other people; are more likely to show blunted empathy toward others; are less likely to direct attention to the eyes of attachment figures; and adapt less well to changes in reward-punishment contingencies.

Functional magnetic resonance imaging (fMRI) studies of children exhibiting disruptive behaviors and CU traits have largely focused on probing the functioning of brain areas implicated in affective/reward processing (see Fig. 1)[†]. These studies have indicated a neural activity profile consistent with low emotional

[†] The notes appear after the main text.




Fig. 1. Neural circuitry involved in emotion, reward, and empathic processing; implicated in studies of children and elevated risk of developing psychopathy. Clearly, most of brain's emotional/empathic processes are achieved by functional integration across several brain areas, that is, although broad functions for these brain areas are discussed in this figure legend (below), many of these areas work together to achieve appropriate behavioral outcome for the organism. For example, amygdala, together with prefrontal cortex (PFC) and ventral striatum, form a network of structures involved in processing the current value of stimuli and various PFC/anterior cingulate cortex (ACC) regions are either directly or indirectly connected with amygdala to achieve emotion regulation via different mechanisms, such as reappraisal. *The amygdala* is a subcortical region that is important for processing the current value of stimuli. The amygdala has a critical role in several affective processes, such as mediating conditioned emotional responses, responding to various emotional stimuli (including facial expressions of emotion), and in social behavior toward conspecifics. *The striatum* is a subcortical region that plays a role in modulating behavior toward potentially rewarding stimuli, particularly stimuli that hold a high subjective reward value to an individual. *The ACC* is thought to play a distinct role in complex aspects of emotion, such as processing moral emotions (e.g. guilt), empathy, self-regulation of negative emotions, and action reinforcement (route by which reward history influences action choice). *The anterior insula (AI)* plays an important role in sensory integration and interoceptive awareness and may be involved in awareness of unpleasant feelings during empathy for pain. *PFC*: Various sectors of the PFC have been implicated in emotion. The areas that have received most attention in affective neuroscience studies of children at risk for developing psychopathy include the orbitofrontal cortex (OFC); ventromedial PFC (vmPFC); ventrolateral PFC (vLPFC); and dorsolateral PFC (DLPFC). The OFC is thought to implement rapid stimulus–reinforcement associations and the correction of these associations when the contingencies of reinforcement change, while the vmPFC is thought to represent the elementary positive and negative affective states in the absence of immediately present incentives. A more rostral region of the MPFC has been implicated in the processing of more complex social emotions such as guilt and embarrassment. Current research suggests that vLPFC integrates affective information and supports response selection by increasing the salience of alternative motor response option representations through interactions with the striatum. It is also associated with effortful regulation of negative affect, via connections with subcortical structures including the amygdala. DLPFC, in turn, is thought to increase attentional control of task-relevant stimulus features and represent goal states toward which more elementary positive and negative affective states are directed [Fig. 1 is reproduced with permission from Viding & McCrory (2015)].

responsiveness to others' distress/pain and poor ability to learn from reinforcement information (Blair *et al.* 2014). Reduced amygdala activity to fearful face in children exhibiting disruptive behaviors with high levels of CU traits relative to typically developing children, children with attention-deficit/hyperactivity disorder (ADHD), and children exhibiting disruptive behaviors without CU traits has been reported in several studies (see Marsh *et al.* 2008; Jones *et al.* 2009; Viding *et al.* 2012; Blair *et al.* 2014), and one study found that the association between CU traits and proactive aggression is partially mediated by low amygdala reactivity to fearful faces (Lozier *et al.* 2014). Cohn *et al.* studied fear conditioning in boys with

early offending histories and reported that CU traits were negatively associated with anterior cingulate cortex activity during fear conditioning (Cohn *et al.* 2013). Using an attention-to-emotion paradigm, White *et al.* (2012a) found atypically low amygdala activation to fear stimuli in children with disruptive behaviors and CU traits under low attentional load conditions; fear stimuli typically elicit amygdala activation in healthy volunteers (White *et al.* 2012a). Furthermore, attentional cueing by eye-gaze of a fearful face does not seem to elicit activation in the dorsal frontoparietal endogenous attention-orienting network in children with disruptive behaviors and high CU traits, although this network is reliably activated in

typically developing children under such conditions (White *et al.* 2012b).

Reduced amygdala and insula activity in children with disruptive behaviors and high CU traits are also seen when these children engage in more complex forms of social judgment regarding other people's distress, such as categorization of legal and illegal behaviors in moral judgment tasks (Marsh *et al.* 2011), making decisions about appropriate responses to the distress of others (Sebastian *et al.* 2012), or making decisions about whether to benefit self by harming others (Sakai *et al.* 2017). Finally, five recent studies of children exhibiting disruptive behaviors and varying levels of CU traits (four involving fMRI, and the other brain event-related potential measurement) have reported atypical neural reactivity to other people's pain (Cheng *et al.* 2012; Lockwood *et al.* 2013; Marsh *et al.* 2013; Michalska *et al.* 2016; Yoder *et al.* 2016). Collectively, these studies implicate reduced activity and altered connectivity in children with disruptive behaviors and high levels of CU traits in a network of brain areas known to be associated with empathy for other people's pain in healthy individuals. This network encompasses a number of brain regions including the anterior insula, posterior insula, anterior cingulate cortex, and the amygdala. Importantly, this profile of reduced neural reactivity to expressions of pain is not coupled with difficulty in understanding intentionality on the part of others (Cheng *et al.* 2012).

Atypical ventromedial prefrontal cortex (vmPFC) and orbitofrontal cortex (OFC) response to punishment and reward in children with disruptive behaviors and CU traits has also been reported (Finger *et al.* 2008, 2011). For example, Finger *et al.* (2008) reported that both typically developing children and children with ADHD showed a reduction in vmPFC activity following an unexpected punishment. Such reduction in vmPFC activity has been shown to co-occur with prediction error, that is, when an actual outcome differs from the expected outcome (Mitchell, 2011). In contrast, youth with disruptive behaviors and CU traits did not show this reduction in vmPFC activity. In another study that used a passive avoidance task in which participants had to learn which stimuli were 'good' (rewarded) and which were 'bad' (unrewarded), Finger *et al.* (2011) found that children with disruptive behaviors and CU traits showed less OFC and caudate responsiveness to early stimulus-reinforcement exposure, and less OFC responsiveness to rewards. These neural differences can be interpreted as reflecting compromised sensitivity to early reinforcement information in the vmPFC/OFC and caudate, and compromised sensitivity to reward outcome information in the vmPFC/OFC, in adolescents with disruptive behaviors and CU traits. Subsequent

work, however, suggests that the difficulties in reinforcement learning may not be unique to disruptive behaviors with CU traits, but may instead be a common problem among children with disruptive behaviors as a whole (White *et al.* 2013, 2014) and relate to the levels of conduct disorder symptoms (White *et al.* 2016a, b).

Collectively, these functional neuroimaging findings are largely in line with those typically reported in studies of psychopathic adults suggesting that assessing disruptive behaviors and CU traits identifies children who present with similar risk markers to those seen in adults with psychopathy (e.g. Kiehl *et al.* 2001; Birbaumer *et al.* 2005; for a comprehensive review, see Seara-Cardoso & Viding, 2014). These findings also suggest functional neural bases for why these children respond less to other people's distress and often make and repeat disadvantageous decisions. What is not clear from the current (cross-sectional) evidence base is the degree to which neurocognitive biases associated psychopathic features in children can change. Is amelioration in neurocognitive differences associated with behavioral improvement in some children? Or do these children achieve behavioral improvement via other, compensatory mechanisms? Answering questions like this will also constrain how 'biomarkers' such as neuroimaging data may be used. Neuroimaging 'biomarkers' may be primarily useful in informing how treatments might be best constrained (i.e. should you try to recalibrate affective processing or should you deploy approaches that focus on compensatory mechanisms) or they may, in some cases, provide additional risk prediction. Longitudinal data are required to answer these questions.

Findings from genetically informative studies

In interpreting findings from genetic/epigenetic studies, it is of critical importance to keep in mind that *there are no genes that directly give rise to the development of psychopathy*. Genes code for proteins that influence characteristics such as neurocognitive vulnerabilities that may in turn increase the *risk for* developing psychopathic features. The neurocognitive vulnerabilities associated with disruptive behaviors that occur together with CU traits are at least partially distinct from those associated with disruptive behaviors more broadly (see 'Findings from neurocognitive studies' section). This suggests that the risk alleles for psychopathy may be in part independent from the risk alleles for disruptive behaviors more broadly. Although an individual's genome likely limits a 'range for phenotypic expression', it does not pre-specify how an individual will turn out. The specific developmental trajectory of any individual is determined by a

complex interplay between genetic propensities and other factors that constrain how those genetic propensities are expressed at several different levels of analysis. Genetic variants that are implicated as risk genes for psychopathic presentation may include several genes that confer advantages, as well as disadvantages, depending on the environmental context. It is also of critical importance to remember that an individual's genetic predisposition influences the types of environments the individual will encounter, a phenomenon known as gene–environment correlation (rGE). For example, an individual's genetic predisposition may influence behaviors that will in turn evoke different reactions in others. In order to understand how environmental risk operates in relation to the development of psychopathy, it is therefore important to conduct research into environmental risk within genetically informative study designs.

Here we review genetically informative findings in relation to CU traits, with particular emphasis on findings in the context of disruptive behaviors. This is to allow comparability with the neurocognitive literature of developmental risk for psychopathy that has primarily focused on differentiating children with disruptive behavior on CU traits. Individual differences in CU are typically estimated to be moderately to strongly heritable using the standard twin design, which compares resemblance in monozygotic (MZ) twins and dizygotic twins, in community samples of children and adolescents (range of heritability estimates =0.45–0.72; see Viding & McCrory, 2012; Ficks *et al.* 2014; Flom & Saudino, 2017; see also Tuvblad *et al.* 2017 for a recent study with a lower heritability estimate of 0.25 in 5-year-old twins). It is also of interest to note that disruptive behaviors accompanied by high levels of CU traits appear strongly heritable, whereas disruptive behaviors accompanied by low levels of CU traits appear to be more strongly influenced by environmental factors (Viding *et al.* 2005, 2008).

Twin studies suggest that there is considerable overlap in the genes that influence CU traits and conduct/externalizing problems, but that there are also unique genetic influences on CU (Viding *et al.* 2007; Forsman *et al.* 2008; Bezdjian *et al.* 2011). This finding is consistent with evidence indicating that high levels of CU can occur in the absence of clinical levels of disruptive behaviors (Frick *et al.* 2003; Kumsta *et al.* 2012), although this is rare. Persistent high levels of CU typically occur in the presence of disruptive behaviors (Fontaine *et al.* 2011). Twin research findings also suggest that observed stability in CU/psychopathic traits is largely driven by genetic influences (Forsman *et al.* 2008; Fontaine *et al.* 2010; Flom & Saudino, 2017).

Only a handful of candidate gene-association studies to date have focused on CU traits in *children or*

adolescents with most of these studies focusing on samples with some disruptive behavior. These studies have tentatively implicated variants of genes related to the serotonin and oxytocin systems (e.g. Fowler *et al.* 2009; Beitchman *et al.* 2012; Malik *et al.* 2012; Dadds *et al.* 2013; Moul *et al.* 2013). Findings from these candidate gene studies need to be investigated in larger samples to evaluate whether they reflect true replicable associations; however, selecting candidate genes is not straightforward and can lead to unadjusted multiple testing. Because genetic risk may in many cases only 'penetrate' in the presence of environmental risk, genetic studies should carefully document the environmental risk factors in their samples to increase interpretability of the findings, and thereby enhance our understanding of how genetic risk translates to disorder outcomes. For example, one interesting study reported that the long allele of a serotonin transporter polymorphism, found to be related to low amygdala reactivity in prior research, was only a risk genotype for elevated CU traits in the context of lower socioeconomic status (SES) backgrounds (Sadeh *et al.* 2010). This finding suggests that the degree to which genetic vulnerability to CU traits is expressed is dependent on the degree to which a child is exposed to environmental disadvantage. However, further studies are required to also investigate the significance of this finding in relation to developing psychopathy in adulthood.

The association studies highlighted above have only considered a limited number of candidate genes. However, an increasing number of genome-wide association (GWA) studies are now being published in the field of psychiatric genetics. These studies systematically scan the genome with hundreds of thousands of DNA markers, made possible by DNA arrays. GWA studies for psychiatric phenotypes have shown that genome-wide 'hits' are often in genes not previously hypothesized to influence the phenotype, or in parts of the genome other than genes themselves (Visscher *et al.* 2012). GWA studies focusing specifically on CU traits (both alone and in combination with disruptive behaviors) suggest that much larger samples will be needed to detect novel associations with common single nucleotide polymorphisms that account for far <1% of the variance (Viding *et al.* 2010, 2013). Furthermore, a genome-wide complex trait analysis study suggests that most of the genetic variance that is important for explaining genetically driven individual differences in CU traits or disruptive behaviors is not due to the additive effects of common genetic variants (Trzaskowski *et al.* 2013). This means that the search for genetic influences on psychopathy is likely to be complicated by the presence of gene–gene interactions and rare variants, as well as gene–environment

interplay. As such, it is questionable how reliable the associations found for individual candidate genes will prove to be. Methods to identify gene–gene and gene–environment interactions are required, as is whole-genome sequencing to detect rare variants that might contribute to heritability of psychopathic presentation, but which would not be detected by candidate gene or GWAS methods. It may be possible to use information from such studies to indicate the most appropriate preventative approaches in the future, as well as inform how genetic features may constrain environmental inputs. Research into genetic contributions to psychopathic features, including studies using novel epigenetic approaches that may also help uncover mechanisms of gene–environment interplay, is likely to progress greatly in the coming decade. For example, two recent epigenetic studies have found that higher CU traits are associated with greater methylation of the oxytocin receptor gene (Cecil *et al.* 2014; Dadds *et al.* 2014), although it remains unclear whether this methylation pattern indexes environmentally or genetically driven epigenetic processes.

Harsh and negative parenting has been associated with higher levels of CU traits and disruptive behaviors, while a warm parental style has been associated with lower levels of CU traits and disruptive behaviors in children (for a review, see Waller *et al.* 2013), but it is not self-evident that such parenting correlates reflect purely environmental causal influences of parenting on behavioral development. Parents with genetic risk factors for psychopathy/antisocial behavior are likely to display parenting behaviors in line with these risks (e.g. harsh parenting) and may also pass on some risk genes to their offspring, which can increase the chance of developing disruptive behaviors and lack of empathy. This means that part of the association between less harsh parenting and CU traits/disruptive behaviors may represent a genetic confound (an example of passive rGE). We also know that children with psychopathic features are extremely challenging to a parent. Given that they typically show diminished empathy for others, less remorse, a tendency to manipulate others, self-serving behaviors, impulsivity, and little interest in seeking the approval of adults, it is not difficult to imagine that common parenting strategies would be quickly exhausted. It is therefore likely that children with psychopathic features evoke different parenting reactions than less challenging children and research suggests that this is the case (Hawes *et al.* 2011) (an example of evocative rGE).

To date only three genetically informative longitudinal studies have investigated parenting and development of psychopathic features, all focusing on CU traits (Viding *et al.* 2009; Hyde *et al.* 2016; Waller *et al.* 2016). Results from the first of these studies,

capitalizing on a MZ-twin differences design, suggest that the association between harsh and negative parenting and higher levels of CU traits in children may, at least in part, reflect genetic vulnerability within families (Viding *et al.* 2009). This could either reflect a shared genetic vulnerability for poor parenting and CU temperament, or an effect of CU temperament in evoking negative/harsh parenting. Complementing this work, two recent studies using the same adoption cohort found that level of antisocial behavior and fearlessness in the *biological mother* predicted early CU behaviors in toddlers who had been adopted. However, high levels of *adoptive mother* positive reinforcement were able to buffer the effects of heritable risk for CU behaviors (Hyde *et al.* 2016; Waller *et al.* 2016). These findings are extremely encouraging, although it is important to bear in mind that parents in adoptive families are typically very motivated to undertake the challenges of parenting and are also often relatively well-resourced. By contrast, in biological families, parents of children with CU traits are more likely to have a host of genetic and contextual risk factors (including socio-economic disadvantage), which can pose challenges for promoting interventions that seek to increase positive reinforcement behaviors toward the child – particularly if that child is challenging.

A few studies to date have also reported an association between CU traits and disorganized attachment in children with disruptive behaviors (Pasalich *et al.* 2011; Bohlin *et al.* 2012). However, no genetically informative longitudinal studies exist that could elucidate the degree to which CU tendencies may drive the development of disorganized attachment in at-risk children. Several studies by Dadds *et al.* (2011, 2012, 2014) suggest that compared with other children with disruptive behaviors, those with high levels of CU traits make less eye contact with their mothers in both free play and directed situations. The mothers of children with high levels of CU traits, on the other hand, do not differ from mothers of non-CU children with disruptive behaviors in the amount of eye contact they attempt with their children. This suggests that children with high CU traits and disruptive behaviors bring a number of challenges to the parent–child relationship, which may influence the formation of attachment style. It will be important to identify within a longitudinal design the features of children with high CU traits and disruptive behaviors that present a challenge to dyadic interactions, as well as carefully characterize the degree to which these are shared with other children with problematic presentation *v.* unique to those at risk for developing psychopathy. Other factors, apart from parenting and parental attachment, are also likely to contribute to the development of

the combination of disruptive behaviors and CU traits and warrant mention. These include SES, peer relationships, cognitive ability, and pre- and peri-natal risk factors (Viding & McCrory, 2015). For example, peer relationships of children with high levels of CU traits are characterized by less stability and greater conflict (Muñoz *et al.* 2008), and children with high levels of CU traits associate more frequently with delinquent friends (Kimonis *et al.* 2004). However, the extant evidence base is not able to elucidate the degree to which these associations are causal or correlational (e.g. related to the phenomenon of rGE). Interestingly, delinquent children with CU traits do not seem to be influenced by their friends' peer delinquency (Kerr *et al.* 2012); rather, their delinquent behavior appears to strongly influence their friends' delinquency (Kerr *et al.* 2012).

The current evidence base clearly indicates that both genetic and environmental risk factors contribute to developmental risk for psychopathy. We are at present some way away from identifying the precise genes that represent the genetic risk. The extant data suggest that finding such genes will be complicated by the presence of gene-gene interactions, as well as rare genetic variants. Findings to date also indicate that although family environmental factors that are associated with disruptive behaviors and CU traits may in part reflect genetic endowment within families, there is also manifest potential for positive environmental factors – such as warm and positive parenting – to ameliorate the development of psychopathic presentation.

Future research needs

Although substantive progress has been made in our understanding of the neurocognitive functioning of individuals at risk of developing psychopathy, as well as the origins of that risk, there are still a number of important questions to be addressed. First, only a limited range of tasks have been used in neurocognitive research (either experimental or imaging studies), the sample sizes in the extant studies (including our own) have often been modest, and the generalizability of the neurocognitive findings across different groups of people has not been systematically investigated. While the current body of findings contains many conceptual replications, exact replications are rare. There are some seemingly conflicting findings and it is not currently clear to what extent these reflect task, trait measurement, or sample differences (see, e.g. Hyde *et al.* 2016; Martin-Key *et al.* 2016). It is important for future studies to use well-constructed measures that are able to clearly isolate specific affective/cognitive processes of interest, and to study whether the findings hold for both sexes or across cultures or ethnicities. It is

also important to systematically investigate the aspects of developmental risk for psychopathy beyond CU traits, for example, those neurocognitive factors associated with seemingly impulsive and maladaptive decision-making and which have recently been isolated in adults with psychopathy (e.g. Baskin-Sommers *et al.* 2016; Hosking *et al.* 2017).

Second, there are no longitudinal studies with neuroimaging measures that shed light on *how* specific processing biases related to psychopathy risk develop nor how malleable such processing biases might be. We also do not know whether clinical improvement is associated with alterations in the neurocognitive systems associated with the risk of psychopathy or whether symptom changes relate to altered neural processing in other brain regions that may reflect 'compensatory' mechanisms.

A related, third point, concerns the lack of longitudinal data on neurocognitive functioning embedded within a rich set of other measures – including genetic and epigenetic data, and data on environmental risk and protective factors. As we have described, the specific developmental trajectory of any individual is determined by a complex interplay between genetic propensities and contextual factors that constrain how those genetic propensities are expressed throughout different developmental stages. Longitudinal work incorporating different levels of analysis would enable researchers to investigate a novel set of questions. For example, it could shed light on how genetic vulnerability might bias affective processing and how such a bias might be further potentiated and canalized given certain environmental contingencies *or* how affective processing biases themselves (in both children and their parents) may shape the environment. There are a number of factors that make conducting this kind of research particularly challenging. We have not yet identified a reasonable set of genes that we can confidently say are associated with the risk for developing psychopathy. The best way to probe neurocognitive biases across different ages is also not clear, nor will it be trivial to combine the different levels of analyses from the point of view of securing a sufficient sample size. We also have a poor understanding of how environmental risk operates in relation to the development of psychopathy and which are the most critical factors to focus on at different developmental stages.

Fourth, we need to investigate heterogeneity in the causal processes that may lead to the development of psychopathic features: such features may emerge in different children for different reasons. Originally, Cleckley (1976) described individuals with psychopathy as being low in anxiety and studies of children at risk for psychopathy often report low levels of co-occurring anxiety in these children (e.g. Frick *et al.* 1999). However, we now

know that a subgroup of children who present with CU traits and disruptive behaviors (a combination taken as a proxy for the risk of developing psychopathy) also present with co-occurring anxiety (Kimonis *et al.* 2011, 2012). It appears that these children are also characterized with more frequent experiences of social and environmental adversity (such as maltreatment) than children with CU traits and disruptive behaviors who present without anxiety. Currently, we do not know the degree to which children exposed to adversity who present with CU, disruptive behaviors, and co-morbid anxiety share genetic, neurocognitive, hormonal, and psychophysiological characteristics with children who exhibit CU traits and disruptive behaviors without anxiety. Emerging research suggests that those with co-morbid anxiety may look more like maltreated and anxious children in general (i.e. showing over-reactivity to affective stimuli, rather than the under-reactivity that is typical for non-anxious high-CU children with disruptive behaviors; Kimonis *et al.* 2012, 2016, 2017). In other words, it is plausible that there may be both genetically and environmentally 'weighted' pathways to psychopathic presentation, and one epigenetic study is in line with this proposal (Cecil *et al.* 2014). Future research efforts are needed to systematically investigate the neurocognitive presentation and developmental trajectories of children who present with psychopathic features and high *v.* low levels of concurrent anxiety.

Treatment implications

While some children present with stable high levels of CU traits and disruptive behaviors, for others these traits either diminish or increase with age (e.g. Burke *et al.* 2007; Fontaine *et al.* 2010). Encouragingly from a clinical perspective, even for those children and youth who show stable high CU traits and disruptive behaviors over time, interventions can contribute to reductions in disruptive behaviors (Waller *et al.* 2013; Hawes *et al.* 2014; Wilkinson *et al.* 2015). Parent training interventions delivered in early childhood have also been shown to produce lasting reductions in CU traits (see Hawes *et al.* 2014), although treatment gains achieved with older children are sometimes more modest (Bjørnebekk & Kjøbli, 2017). The challenge over the next decade will be to more comprehensively delineate what precisely works for these children and how current intervention and prevention programs can be optimized in ways that improve engagement as well as clinical outcomes.

To optimize existing interventions (as well as engagement), we may need a better understanding of rGE processes. Such research has the potential to inform clinical approaches that promote warm and consistent

parenting. A child with psychopathic features is more likely to evoke negative parenting responses (as well as a sense of parental inadequacy) and will furnish more infrequent occasions for parental praise or reward. This in turn increases the likelihood that the pattern of parent-child interaction becomes largely negative in tone. Furthermore, the biological parents of such a child may themselves share some of the same vulnerabilities that characterize their child and may find it harder to implement many aspects of typical parenting intervention programs. Biological families of children with psychopathic features may need additional (or different forms of) clinical support in order to facilitate a shift toward more positive patterns of parent-child interaction. These could include intensive and supported reward token-economy adjuncts and extensive support in increasing warm parent-child interactions (see Armstrong & Kimonis, 2013 for a promising case study).

In addition, it will be important to examine the degree to which the neurocognitive biases associated with psychopathic features are malleable as this will help inform the development of adjuncts to existing parenting- or family-focused clinical interventions. If it transpires, for example, that a given bias is not malleable, it may be more fruitful to focus on the development of compensatory processes that can have the effect of normalizing behavior. Relatedly, we also need to investigate whether there are particularly sensitive developmental periods during which the neurocognitive functioning may be most responsive to intervention. Given that the children's processing of the world around them is likely to impact upon their social interactions, development of treatment adjuncts can also be seen as providing extra scaffolding for a positive parent-child relationship and attenuating rGE processes that might otherwise unfold over development.

Conclusions

Despite substantial accumulation of new evidence, we still lack a coherent understanding of how genetic and environmental risk factors shape the development of neurocognitive processing biases that increase risk for psychopathy, or how those processing biases themselves may come to shape the individual's environmental exposures over time. Longitudinal research combining multiple levels of analyses is critical to progress the field, but conducting such research will be challenging. It will require systematic development of measures that are able to index the development of specific processing biases from infancy to adulthood, as well identification of age-relevant environmental risk factors and their mode of operation. For example, why is reduced face preference during infancy associated with increased likelihood of CU traits in early

childhood (Bedford *et al.* 2015)? Does it, perhaps combined with a genetic predisposition to a fearless temperament, lead to fewer opportunities to observe a caregiver's frightened face after a loud noise or flashing light? Might that, in turn, lead to a less consolidated representation of what arousal state is associated with a frightened facial expression and would that make it harder for a child to empathize with another person's distress? Does regular positive reinforcement by a parent promote prosocial behavior in ways that help the child integrate socially, attenuating difficulties in empathizing? Currently we do not know, but it will be important to find out. Longitudinal investigations including data from different levels of analyses are either scarce or absent, and we do not currently have good data on how the child's predisposition (variously quantified) contributes to the parent-child or child-peer dynamic over time. Fortunately, a number of research groups are engaged in conducting systematic, innovative research to answer these and other questions. We are confident that we will be able to write a very different, much improved review on the development of psychopathy in 10 years time.

Note

- ¹ For an excellent review of structural brain imaging findings in this area see Rogers & De Brito (2016). Here we only review findings from functional neuroimaging studies.

Acknowledgements

EV and EJM were supported by a UK Medical Research Council grant (MR/ K014080/1) during the writing of this paper. E.V. is a Royal Society Wolfson Research Merit Award holder and a British Academy Mid-Career Fellow.

References

- American Psychiatric Association** (2013). *Diagnostic and Statistical Manual of Mental Disorders*, 5th edn. American Psychiatric Association: Washington, DC.
- Andershed H, Gustafson SB, Kerr M, Stattin H** (2002). The usefulness of self-reported psychopathy-like traits in the study of antisocial behaviour among non-referred adolescents. *European Journal of Personality* **16**, 383–402.
- Armstrong K, Kimonis ER** (2013). Parent-child interaction therapy for the treatment of Asperger's disorder in early childhood: a case study. *Clinical Case Studies* **12**, 60–72.
- Baskin-Sommers A, Stuppy-Sullivan AM, Buckholtz JW** (2016). Psychopathic individuals exhibit but do not avoid regret during counterfactual decision making. *Proceedings of the National Academy of Sciences of the United States of America* **113**, 14438–14443.
- Bedford R, Pickles A, Sharp H, Wright N, Hill J** (2015). Reduced face preference in infancy: a developmental precursor to callous-unemotional traits? *Biological Psychiatry* **78**, 144–150.
- Beitchman JH, Zai CC, Muir K, Berall L, Nowrouzi B, Choi E, Kennedy JL** (2012). Childhood aggression, callous-unemotional traits and oxytocin genes. *European Child & Adolescent Psychiatry* **21**, 125–132.
- Bezdjian S, Raine A, Baker LA, Lynam DR** (2011). Psychopathic personality in children: genetic and environmental contributions. *Psychological Medicine* **41**, 589–600.
- Birbaumer N, Veit R, Lotze M, Erb M, Hermann C, Grodd W, Flor H** (2005). Deficient fear conditioning in psychopathy: a functional magnetic resonance imaging study. *Archives of General Psychiatry* **62**, 799–805.
- Bjornebekk G, Kjøbli J** (2017). Observed callousness as a predictor of treatment outcomes in parent management training. *Clinical Child Psychology and Psychiatry* **22**, 59–73.
- Blair RJR, Leibenluft E, Pine DS** (2014). Conduct disorder and callous-unemotional traits in youth. *New England Journal of Medicine* **371**, 2207–2216.
- Bohlin G, Eninger L, Brocki KC, Thorell LB** (2012). Disorganized attachment and inhibitory capacity: predicting externalizing problem behaviors. *Journal of Abnormal Child Psychology* **40**, 449–458.
- Burke JD, Loeber R, Lahey BB** (2007). Adolescent conduct disorder and interpersonal callousness as predictors of psychopathy in young adults. *Journal of Clinical Child and Adolescent Psychology* **36**, 334–346.
- Cecil CA, Viding E, Barker ED, Guiney J, McCrory EJ** (2014). Double disadvantage: the influence of childhood maltreatment and community violence exposure on adolescent mental health. *Journal of Child Psychology and Psychiatry* **55**, 839–848.
- Cheng Y, Hung AY, Decety J** (2012). Dissociation between affective sharing and emotion understanding in juvenile psychopaths. *Development and Psychopathology* **24**, 623–636.
- Cleckley H** (1976). *The Mask of Sanity*, 1st edn/5th edn. Mosby: St. Louis, MO.
- Cohn MD, Popma A, van den Brink W, Pape LE, Kindt M, van Domburgh L, Doreleijers TAH, Veltman DJ** (2013). Fear conditioning, persistence of disruptive behavior and psychopathic traits: an fMRI study. *Translational Psychiatry* **3**, e319.
- Colins OF, Andershed H, Frogner L, Lopez-Romero L, Veen V, Andershed AK** (2014). A new measure to assess psychopathic personality in children: the child problematic traits inventory. *Journal of Psychopathology and Behavioral Assessment* **36**, 4–21.
- Dadds MR, Allen JL, McGregor K, Woolgar M, Viding E, Scott S** (2014). Callous-unemotional traits in children and mechanisms of impaired eye contact during expressions of love: a treatment target? *Journal of Child Psychology and Psychiatry* **55**, 71–780.
- Dadds MR, Allen JL, Oliver BR, Faulkner N, Legge K, Moul C, Woolgar M, Scott S** (2012). Love, eye contact and the

- developmental origins of empathy *v.* psychopathy. *The British Journal of Psychiatry* **200**, 191–196.
- Dadds MR, Jambrak J, Pasalich D, Hawes DJ, Brennan J** (2011). Impaired attention to the eyes of attachment figures and the developmental origins of psychopathy. *Journal of Child Psychology and Psychiatry* **52**, 238–245.
- Dadds MR, Moul C, Cauchi A, Hawes DJ, Brennan J** (2013). Replication of a ROBO2 polymorphism associated with conduct problems but not psychopathic tendencies in children. *Psychiatric Genetics* **23**, 251–254.
- de Wied M, van Boxtel A, Matthys W, Meeus W** (2012). Verbal, facial and autonomic responses to empathy-eliciting film clips by disruptive male adolescents with high *versus* low callous-unemotional traits. *Journal of Abnormal Child Psychology* **40**, 211–223.
- Essau CA, Sasagawa S, Frick PJ** (2006). Callous-unemotional traits in a community sample of adolescents. *Assessment* **13**, 454–469.
- Ficks CA, Dong L, Waldman ID** (2014). Sex differences in the etiology of psychopathic traits in youth. *Journal of Abnormal Psychology* **123**, 406–411.
- Finger EC, Marsh AA, Blair KS, Reid ME, Sims C, Ng P, Pine DS, Blair RJR** (2011). Disrupted reinforcement signaling in the orbitofrontal cortex and caudate in youths with conduct disorder or oppositional defiant disorder and a high level of psychopathic traits. *American Journal of Psychiatry* **168**, 152–162.
- Finger EC, Marsh AA, Mitchell DG, Reid ME, Sims C, Budhani S, Kosson DS, Chen G, Towbin KE, Leibenluft E, Blair JR** (2008). Abnormal ventromedial prefrontal cortex function in children with psychopathic traits during reversal learning. *Archives of General Psychiatry* **65**, 586–594.
- Flom M, Saudino KJ** (2017). Callous–unemotional behaviors in early childhood: genetic and environmental contributions to stability and change. *Development and Psychopathology* **29**, 1227–1234.
- Fontaine NM, McCrory EJ, Boivin M, Moffitt TE, Viding E** (2011). Predictors and outcomes of joint trajectories of callous–unemotional traits and conduct problems in childhood. *Journal of Abnormal Psychology* **120**, 730–742.
- Fontaine NM, Rijdsdijk FV, McCrory EJ, Viding E** (2010). Etiology of different developmental trajectories of callous-unemotional traits. *Journal of the American Academy of Child & Adolescent Psychiatry* **49**, 656–664.
- Forsman M, Lichtenstein P, Andershed H, Larsson H** (2008). Genetic effects explain the stability of psychopathic personality from mid-to late adolescence. *Journal of Abnormal Psychology* **117**, 606–617.
- Fowler T, Langley K, Rice F, van den Bree MB, Ross K, Wilkinson LS, Owen MJ, O'donovan MC, Thapar A** (2009). Psychopathy trait scores in adolescents with childhood ADHD: the contribution of genotypes affecting MAOA, 5HTT and COMT activity. *Psychiatric Genetics* **19**, 312–319.
- Frick PJ, Cornell AH, Barry CT, Bodin SD, Dane HE** (2003). Callous-unemotional traits and conduct problems in the prediction of conduct problem severity, aggression, and self-report of delinquency. *Journal of Abnormal Child Psychology* **31**, 457–470.
- Frick PJ, Hare RD** (2001). *The Antisocial Process Screening Device*. Multi-Health Systems: Toronto, ON.
- Frick PJ, Lilienfeld SO, Ellis M, Loney B, Silverthorn P** (1999). The association between anxiety and psychopathy dimensions in children. *Journal of Abnormal Child Psychology* **27**, 383–392.
- Frick PJ, Ray JV, Thornton LC, Kahn RE** (2014). Annual research review: a developmental psychopathology approach to understanding callous-unemotional traits in children and adolescents with serious conduct problems. *Journal of Child Psychology and Psychiatry* **55**, 532–548.
- Frogner L, Gibson CL, Andershed AK, Andershed H** (2016). Childhood psychopathic personality and callous-unemotional traits in the prediction of conduct problems. *American Journal of Orthopsychiatry* Oct 27, Epub ahead of print.
- Hare RD** (2003). *Manual for the Revised Psychopathy Checklist*, 2nd edn. Multi-Health Systems: Toronto, ON.
- Hare RD, Neumann CS** (2008). Psychopathy as a clinical and empirical construct. *Annual Review of Clinical Psychology* **4**, 217–246.
- Hawes DJ, Dadds MR, Frost AD, Hasking PA** (2011). Do childhood callous-unemotional traits drive change in parenting practices? *Journal of Clinical Child & Adolescent Psychology* **40**, 507–518.
- Hawes DJ, Price MJ, Dadds MR** (2014). Callous-unemotional traits and the treatment of conduct problems in childhood and adolescence: a comprehensive review. *Clinical Child and Family Psychology Review* **17**, 248–276.
- Hodsoll S, Lavie N, Viding E** (2014). Emotional attentional capture in children with conduct problems: the role of callous-unemotional traits. *Frontiers in Human Neuroscience* **8**, 570.
- Hosking JG, Kastman EK, Dorfman HM, Samanez-Larkin GR, Baskin-Sommers A, Kiehl KA, Newman JP, Buckholz JW** (2017). Disrupted prefrontal regulation of striatal subjective value signals in psychopathy. *Neuron* **95**, 221–231.
- Hyde LW, Waller R, Trentacosta CJ, Shaw DS, Neiderhiser JM, Ganiban JM, Reiss D, Leve LD** (2016). Heritable and nonheritable pathways to early callous-unemotional behaviors. *American Journal of Psychiatry* **173**, 903–910.
- Jones AP, Happé FG, Gilbert F, Burnett S, Viding E** (2010). Feeling, caring, knowing: different types of empathy deficit in boys with psychopathic tendencies and autism spectrum disorder. *Journal of Child Psychology and Psychiatry* **51**, 1188–1197.
- Jones AP, Riley RD, Williamson PR, Whitehead A** (2009). Meta-analysis of individual patient data *versus* aggregate data from longitudinal clinical trials. *Clinical Trials* **6**, 16–27.
- Kerr M, Van Zalk M, Stattin H** (2012). Psychopathic traits moderate peer influence on adolescent delinquency. *Journal of Child Psychology and Psychiatry* **53**, 826–835.
- Kiehl KA** (2015). *The Psychopath Whisperer: The Science of Those Without Conscience*. Crown: New York.
- Kiehl KA, Smith AM, Hare RD, Mendrek A, Forster BB, Brink J, Liddle PF** (2001). Limbic abnormalities in affective processing by criminal psychopaths as revealed by functional magnetic resonance imaging. *Biological Psychiatry* **50**, 677–684.

- Kimonis ER, Frick PJ, Barry CT** (2004). Callous-unemotional traits and delinquent peer affiliation. *Journal of Consulting and Clinical Psychology* **72**, 956–966.
- Kimonis ER, Frick PJ, Cauffman E, Goldweber A, Skeem J** (2012). Primary and secondary variants of juvenile psychopathy differ in emotional processing. *Development and Psychopathology* **24**, 1091–1103.
- Kimonis ER, Frick PJ, Munoz LC, Aucoin KJ** (2008). Callous-unemotional traits and the emotional processing of distress cues in detained boys: testing the moderating role of aggression, exposure to community violence, and histories of abuse. *Development and Psychopathology* **20**, 569–589.
- Kimonis ER, Goulter N, Hawes DJ, Wilbur RR, Groer MW** (2017). Neuroendocrine factors distinguish juvenile psychopathy variants. *Developmental Psychobiology* **59**, 161–173.
- Kimonis ER, Kennealy PJ, Goulter N** (2016). Does the self-report inventory of callous-unemotional traits predict recidivism? *Psychological Assessment* **28**, 1616–1624.
- Kimonis ER, Skeem J, Cauffman E, Dmitrieva J** (2011). Are secondary variants of ‘juvenile psychopathy’ more reactively violent and less psychosocially mature than primary variants? *Law and Human Behavior* **35**, 381–391.
- Kumsta R, Sonuga-Barke E, Rutter M** (2012). Adolescent callous-unemotional traits and conduct disorder in adoptees exposed to severe early deprivation. *The British Journal of Psychiatry* **200**, 197–201.
- Lilienfeld SO, Andrews BP** (1996). Development and preliminary validation of a self-report measure of psychopathic personality traits in noncriminal population. *Journal of Personality Assessment* **66**, 488–524.
- Lockwood PL, Sebastian CL, McCrory EJ, Hyde ZH, Gu X, De Brito SA, Viding E** (2013). Association of callous traits with reduced neural response to others’ pain in children with conduct problems. *Current Biology* **23**, 901–905.
- Lozier LM, Cardinale EM, VanMeter JW, Marsh AA** (2014). Mediation of the relationship between callous-unemotional traits and proactive aggression by amygdala response to fear among children with conduct problems. *JAMA Psychiatry* **71**, 627–636.
- Lynam DR, Caspi A, Moffitt TE, Loeber R, Stouthamer-Loeber M** (2007). Longitudinal evidence that psychopathy scores in early adolescence predict adult psychopathy. *Journal of Abnormal Psychology* **116**, 155.
- Malik AI, Zai CC, Abu Z, Nowrouzi B, Beitchman JH** (2012). The role of oxytocin and oxytocin receptor gene variants in childhood-onset aggression. *Genes, Brain and Behavior* **11**, 545–551.
- Marsh AA, Blair RJR** (2008). Deficits in facial affect recognition among antisocial populations: a meta-analysis. *Neuroscience & Biobehavioral Reviews* **32**, 454–465.
- Marsh A, Finger E, Mitchell D, Reid M, Sims C, Kosson D, Towbin KE, Leibenluft E, Pine DS, Blair R** (2008). Reduced amygdala response to fearful expressions in children and adolescents with callous-unemotional traits and disruptive behavior disorders. *American Journal of Psychiatry* **165**, 712–720.
- Marsh AA, Finger EC, Fowler KA, Adalio CJ, Jurkowitz IT, Schechter JC, Pine DS, Decety J, Blair RJR** (2013). Empathic responsiveness in amygdala and anterior cingulate cortex in youths with psychopathic traits. *Journal of Child Psychology and Psychiatry* **54**, 900–910.
- Marsh AA, Finger EC, Schechter JC, Jurkowitz IT, Reid ME, Blair RJR** (2011). Adolescents with psychopathic traits report reductions in physiological responses to fear. *Journal of Child Psychology and Psychiatry* **52**, 834–841.
- Martin-Key N, Brown T, Fairchild G** (2016). Empathic accuracy in male adolescents with conduct disorder and higher versus lower levels of callous-unemotional traits. *Journal of Abnormal Child Psychology* 1–13.
- Michalska KJ, Zeffiro TA, Decety J** (2016). Brain response to viewing others being harmed in children with conduct disorder symptoms. *Journal of Child Psychology and Psychiatry* **57**, 510–519.
- Mitchell KJ** (2011). The miswired brain: making connections from neurodevelopment to psychopathology. *BMC Biology* **9**, 23.
- Moul C, Dobson-Stone C, Brennan J, Hawes D, Dadds M** (2013). An exploration of the serotonin system in antisocial boys with high levels of callous-unemotional traits. *PLoS ONE* **8**, e56619.
- Muñoz LC, Kerr M, Besic N** (2008). The peer relationships of youths with psychopathic personality traits a matter of perspective. *Criminal Justice and Behavior* **35**, 212–227.
- Pasalich DS, Dadds MR, Hawes DJ, Brennan J** (2011). Do callous-unemotional traits moderate the relative importance of parental coercion versus warmth in child conduct problems? An observational study. *Journal of Child Psychology and Psychiatry* **52**, 1308–1315.
- Patrick CJ** (2010). Operationalizing the triarchic conceptualization of psychopathy: Preliminary description of brief scales for assessment of boldness, meanness, and disinhibition. *Unpublished test manual, Florida State University, Tallahassee, Florida*. Test retrieved from (<https://www.phenxtoolkit.org/index.php?pageLink=browse.protocoldetails&id=121601>).
- Paulhus DL, Neumann CS, Hare RD** (2016). *Manual for the Self-Report Psychopathy Scale*, 4th edn. Multi-Health Systems: Toronto.
- Rogers JC, De Brito SA** (2016). Cortical and subcortical gray matter volume in youths with conduct problems: a meta-analysis. *JAMA Psychiatry* **73**, 64–72.
- Sadeh N, Javdani S, Jackson JJ, Reynolds EK, Potenza MN, Gelernter J, Lejuez CW, Verona E** (2010). Serotonin transporter gene associations with psychopathic traits in youth vary as a function of socioeconomic resources. *Journal of Abnormal Psychology* **119**, 604–609.
- Sakai JT, Dalwani MS, Mikulich-Gilbertson SK, Raymond K, McWilliams S, Tanabe J, Rojas D, Regner M, Banich MT, Crowley TJ** (2017). Imaging decision about whether to, or not to, benefit self by harming others: adolescents with conduct and substance problems, with or without callous-unemotionality, or developing typically. *Psychiatry Research: Neuroimaging* **263**, 103–112.
- Schwenck C, Mergenthaler J, Keller K, Zech J, Salehi S, Taurines R, Romanos M, Schecklmann M, Schneider W, Warnke A, Freitag CM** (2012). Empathy in children with autism and conduct disorder: group-specific profiles and

- developmental aspects. *Journal of Child Psychology and Psychiatry* **53**, 651–659.
- Seara-Cardoso A, Viding E** (2014). Functional neuroscience of psychopathic personality in adults. *Journal of Personality* **83**, 723–737.
- Sebastian CL, McCrory EJ, Cecil CA, Lockwood PL, De Brito SA, Fontaine NM, Viding E** (2012). Neural responses to affective and cognitive theory of mind in children with conduct problems and varying levels of callous-unemotional traits. *Archives of General Psychiatry* **69**, 814–822.
- Sylvers PD, Brennan PA, Lilienfeld SO** (2011). Psychopathic traits and preattentive threat processing in children: a novel test of the fearlessness hypothesis. *Psychological Science* **22**, 1280–1287.
- Trzaskowski M, Dale PS, Plomin R** (2013). No genetic influence for childhood behavior problems from DNA analysis. *Journal of American Academy of Child and Adolescent Psychiatry* **52**, 1048–1056.
- Tuvblad C, Fanti KA, Andershed H, Colins OF, Larsson H** (2017). Psychopathic personality traits in 5 year old twins: the importance of genetic and shared environmental influences. *European Child & Adolescent Psychiatry* **26**, 469–479.
- Viding E, Blair RJR, Moffitt TE, Plomin R** (2005). Evidence for substantial genetic risk for psychopathy in 7-year-olds. *Journal of Child Psychology and Psychiatry* **46**, 592–597.
- Viding E, Frick PJ, Plomin R** (2007). Aetiology of the relationship between callous-unemotional traits and conduct problems in childhood. *The British Journal of Psychiatry* **190**, s33–s38.
- Viding E, Hanscombe KB, Curtis CJ, Davis OS, Meaburn EL, Plomin R** (2010). In search of genes associated with risk for psychopathic tendencies in children: a two-stage genome-wide association study of pooled DNA. *Journal of Child Psychology and Psychiatry* **51**, 780–788.
- Viding E, Jones AP, Paul JF, Moffitt TE, Plomin R** (2008). Heritability of antisocial behaviour at 9: do callous-unemotional traits matter? *Developmental Science* **11**, 17–22.
- Viding E, McCrory EJ** (2012). Genetic and neurocognitive contributions to the development of psychopathy. *Development and Psychopathology* **24**, 969–983.
- Viding E, McCrory EJ** (2015). Developmental risk for psychopathy. In *Rutter's Child and Adolescent Psychiatry*, 6th edn (ed. A. Thapar, D. S. Pine, J. F. Leckman, S. Scott, M. J. Snowling, E. Taylor), pp. 966–980. John Wiley & Sons.
- Viding E, Price TS, Jaffee SR, Trzaskowski M, Davis OS, Meaburn EL, Haworth CM, Plomin R** (2013). Genetics of callous-unemotional behavior in children. *PLoS ONE* **8**, e65789.
- Viding E, Sebastian CL, Dadds MR, Lockwood PL, Cecil CA, De Brito SA, McCrory EJ** (2012). Amygdala response to preattentive masked fear in children with conduct problems: the role of callous-unemotional traits. *American Journal of Psychiatry* **169**, 1109–1116.
- Viding E, Simmonds E, Petrides KV, Frederickson N** (2009). The contribution of callous-unemotional traits and conduct problems to bullying in early adolescence. *Journal of Child Psychology and Psychiatry* **50**, 471–481.
- Visscher PM, Brown MA, McCarthy MI, Yang J** (2012). Five years of GWAS discovery. *American Journal of Human Genetics* **90**, 7–24.
- Waller R, Dishion TJ, Shaw DS, Gardner F, Wilson MN, Hyde LW** (2016). Does early childhood callous-unemotional behavior uniquely predict behavior problems or callous-unemotional behavior in late childhood? *Developmental Psychology* **52**, 1805–1819.
- Waller R, Gardner F, Hyde LW** (2013). What are the associations between parenting, callous-unemotional traits, and antisocial behavior in youth? A systematic review of evidence. *Clinical Psychology Review* **33**, 593–608.
- White SF, Briggs-Gowan MJ, Voss JL, Petittlerc A, McCarthy KR, Blair RJR, Wakschlag LS** (2016a). Can the fear recognition deficits associated with callous-unemotional traits be identified in early childhood? *Journal of Clinical and Experimental Neuropsychology* **38**, 672–684.
- White SF, Brislin S, Sinclair S, Fowler KA, Pope K, Blair RJR** (2013). The relationship between large cavum septum pellucidum and antisocial behavior, callous-unemotional traits and psychopathy in adolescents. *Journal of Child Psychology and Psychiatry* **54**, 575–581.
- White SF, Fowler KA, Sinclair S, Schechter JC, Majestic CM, Pine DS, Blair RJ** (2014). Disrupted expected value signaling in youth with disruptive behavior disorders to environmental reinforcers. *Journal of American Academy of Child and Adolescent Psychiatry* **53**, 579–588.
- White SF, Marsh AA, Fowler KA, Schechter JC, Adalio C, Pope K, Sinclair S, Pine DS, Blair RJR** (2012a). Reduced amygdala response in youths with disruptive behavior disorders and psychopathic traits: decreased emotional response *versus* increased top-down attention to nonemotional features. *American Journal of Psychiatry* **169**, 750–758.
- White SF, Tyler PM, Erway AK, Botkin ML, Kolli V, Meffert H, Pope K, Blair JR** (2016b). Dysfunctional representation of expected value is associated with reinforcement-based decision-making deficits in adolescents with conduct problems. *Journal of Child Psychology and Psychiatry* **57**, 938–946.
- White SF, Williams WC, Brislin SJ, Sinclair S, Blair KS, Fowler KA, Pine DS, Pope K, Blair JR** (2012b). Reduced activity within the dorsal endogenous orienting of attention network to fearful expressions in youth with disruptive behavior disorders and psychopathic traits. *Development and Psychopathology* **24**, 1105–1116.
- Wilkinson S, Waller R, Viding E** (2015). Practitioner review: involving young people with callous unemotional traits in treatment – does it work? A systematic review. *Journal of Child Psychology and Psychiatry* **57**, 552–565.
- Yoder KJ, Lahey BB, Decety J** (2016). Callous traits in children with and without conduct problems predict reduced connectivity when viewing harm to others. *Scientific Reports* **6**, 20216.